




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 413/06, A61K 31/405	A1	(11) International Publication Number: WO 95/32966 (43) International Publication Date: 7 December 1995 (07.12.95)
(21) International Application Number: PCT/GB95/01249 (22) International Filing Date: 31 May 1995 (31.05.95) (30) Priority Data: 94303928.9 1 June 1994 (01.06.94) EP <i>(34) Countries for which the regional or international application was filed:</i> GB et al. (71) Applicant (for all designated States except US): THE WELL-COME FOUNDATION LIMITED [GB/GB]; Unicom House, 160 Euston Road, London NW1 2BP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): BLADE, Robert, John [GB/GB]; Langley Court, Beckenham, Kent BR3 3BS (GB). PANG, Yih, Sang [GB/GB]; Langley Court, Beckenham, Kent BR3 3BS (GB). SELWOOD, David, Lawrence [GB/GB]; Langley Court, Beckenham, Kent BR3 3BS (GB). (74) Agent: HAMMER, Catriona, MacLeod; The Wellcome Foundation Limited, Langley Court, Beckenham, Kent BR3 3BS (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>

(54) Title: INDOLE DERIVATIVES AS PRODRUGS OF 5-HT₁-LIKE RECEPTOR AGONISTS**(57) Abstract**

A compound of formula (I), wherein A is C₁₋₆alkyl, -O-C₁₋₆alkyl, -O-phenyl or phenyl, optionally substituted by C₁₋₃alkyl or halogen; n is an integer of from 0 to 3; W is a group of formula (i), (ii) or (iii) wherein R is hydrogen or C₁₋₄alkyl, x is -O-, -S-, -NH-, or -CH₂-, Y is oxygen or sulphur and the chiral centre * in formula (i) or (ii) is in its (S) or (R) form or is a mixture thereof in any proportions; and Z is a group of formula (iv) -CH₂CH₂NR¹R², (v) or (vi) wherein R¹ and R² are independently selected from hydrogen and C₁₋₄alkyl and R³ is hydrogen or C₁₋₄alkyl; and salts, solvates and physiologically functional derivatives thereof.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INDOLE DERIVATIVES AS PRODRUGS OF 5-HT₁-LIKE RECEPTOR AGONISTS

The present invention is concerned with new chemical compounds, their preparation, pharmaceutical formulations containing them and their use in medicine, particularly the prophylaxis and treatment of migraine.


Receptors which mediate the actions of 5-hydroxytryptamine (5-HT) have been identified in mammals in both the periphery and the brain. Currently, as many as seven 5-HT receptor classes are proposed (Humphrey et al., Trends Pharmac Sci., 14, 233-236, 1993), although only the classes nominated 5-HT₁, 5-HT₂, 5-HT₃, and 5-HT₄ have established physiological roles. European Patent Specification 0313397 describes a class of 5-HT agonists which act selectively at a particular subtype of 5-HT₁ receptor and are effective therapeutic agents for the treatment of clinical conditions in which a selective agonist for this type of receptor is indicated. For example, the receptor in question mediates selective cranial arterial vasoconstriction and inhibition of plasma protein extravasation into the dura mater evoked by activation of the Vth (trigeminal) nerve. The compounds described in the European specification are therefore beneficial in the treatment or prophylaxis of conditions wherein these actions are indicated, for example, migraine, a condition associated with and/or neurogenically-evoked inflammation dilation of the cranial vasculature. However, it is within the scope of the earlier application that the target tissue may be any tissue wherein action is mediated by 5-HT₁ receptors of the type referred to above.

EP-A-0486666 discloses a class of compounds having exceptional activity at the 5-HT₁ receptor mentioned above and excellent absorption following oral dosing. These properties render the compounds particularly useful for certain medical applications, notably the prophylaxis and treatment of migraine, cluster headache and headache associated with vascular disorders, hereinafter referred to collectively as "migraine".

There has now been discovered a class of compounds which, although having relatively little activity themselves at the 5-HT₁ receptor, act as prodrugs releasing active compound after administration. Such compounds thereby provide active compound with improved metabolic stability and bioavailability.

Thus, according to a first aspect of the present invention there is provided a compound of formula (I):

-2-




wherein


A is C₁₋₆ alkyl, -O-C₁₋₆ alkyl, or -O-phenyl, or phenyl, optionally substituted by C₁₋₃ alkyl or halogen;

n is an integer of from 0 to 3;


W is a group of formula (i), (ii) or (iii)



(i)




(ii)




(iii)

wherein R is hydrogen or C₁₋₄ alkyl, X is -O-, -S-, -NH-, or -CH₂-, Y is oxygen or sulphur and the chiral centre * in formula (i) or (ii) is in its (S) or (R) form or is a mixture thereof in any proportions; and


Z is a group of formula (iv), (v) or (vi)



(iv)



(v)




(vi)

wherein R¹ and R² are independently selected from hydrogen and C₁₋₄ alkyl and R³ is hydrogen or C₁₋₄ alkyl;

and salts, solvates and physiologically functional derivatives thereof.

Preferably W is a group of formula (i) and Z is a group of formula (iv)

In another aspect the present invention provides a compound of the formula (Ia)



wherein

A is C₁₋₆ alkyl, -O-C₁₋₆ alkyl, -O-phenyl or phenyl, optionally substituted by C₁₋₃ alkyl or halogen;

n is an integer from 0 to 3;

R is hydrogen or C₁₋₄ alkyl;

X is -O-, -S-, -NH- or -CH₂-;

Y is oxygen or sulphur; and

R¹ and R² are independently selected from hydrogen and C₁₋₄ alkyl

and salts, solvates and physiologically functional derivatives thereof.

Preferred compounds of formula (Ia) are those wherein n is 1, X is -O- and Y is oxygen.

Compounds wherein R¹ and R² are independently selected from hydrogen and methyl are preferred, with compounds wherein R¹ and R² are both methyl being particularly preferred.

Compounds of formula (I) and (Ia) are capable of existing as optical isomers. All such isomers, individually and as mixtures, are included within the scope of the present invention.

Examples of preferred compounds of the invention include:-

-4-

- 1) 4-[1-Benzoyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate;
- 2) 4-[1-Pivaloyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl] oxazolidin-2-one acetate;
- 3) 4-[1-(2'-toluoyl)-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl] oxazolidin-2-one acetate;
- 4) 4-[1-Benzoyloxycarbonyl-3-[2-(dimethylamino)ethyl] indol-5-ylmethyl]oxazolidin-2-one;
- 5) 4-[1-t-Butyloxycarbonyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl] oxazolidin-2-one acetate; and
- 6) 4-[1-Acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl] oxazolidin-2-one acetate.

Physiologically acceptable salts are particularly suitable for medical applications because of their greater aqueous solubility relative to the parent, ie basic, compounds. Such salts must clearly have a physiologically acceptable anion. Suitable physiologically acceptable salts of the compounds of the present invention include those derived from acetic, hydrochloric, hydrobromic, phosphoric, malic, maleic, fumaric, citric, sulphuric, lactic, or tartaric acid. The succinate and chloride salts are particularly preferred for medical purposes. Salts having a non-physiologically acceptable anion are within the scope of the invention as useful intermediates for the preparation of physiologically acceptable salts and/or for use in non-therapeutic, for example, in vitro, situations.

According to a third aspect of the present invention, there is provided a compound of formula (I) or (Ia) or a physiologically acceptable salt, solvate, or physiologically functional derivative thereof for use as a therapeutic agent, specifically as a "5-HT₁-like" receptor agonist, for example, as a carotid vasoconstrictor or as an inhibitor of neurogenic inflammation in the prophylaxis and treatment of migraine. As indicated, however, target organs for the present compounds other than the carotid vasculature are within the scope of the present invention.

The amount of a compound of formula (I) or (Ia), a salt or solvate thereof, which is required to achieve the desired biological effect will depend on a number of factors such as the specific compound, the use for which it is intended, the means of administration, and

the recipient. A typical daily dose for the treatment of migraine may be expected to lie in the range 0.01 to 5mg per kilogram body weight. Unit doses may contain from 1 to 100mg of a compound of formula (I), for example, ampoules for injection may contain from 1 to 10mg and orally administrable unit dose formulations such as tablets or capsules may contain from 1 to 100mg. Such unit doses may be administered one or more times a day, separately or in multiples thereof. An intravenous dose may be expected to lie in the range 0.01 to 0.15mg/kg and would typically be administered as an infusion of from 0.0003 to 0.15mg per kilogram per minute. Infusion solutions suitable for this purpose may contain from 0.01 to 10mg/ml.

When the active compound is a salt or solvate of a compound of formula (I) the dose is based on the cation (for salts) or the unsolvated compound.

Hereinafter references to "compound(s) of formula (I) or (Ia)" will be understood to include physiologically acceptable salts and solvates thereof.

According to a fourth aspect of the present invention, therefore, there are provided pharmaceutical compositions comprising, as active ingredient, at least one compound of formula (I) or (Ia) and/or a pharmacologically acceptable salt or solvate thereof together with at least one pharmaceutical carrier or excipient. These pharmaceutical compositions may be used in the prophylaxis or treatment of clinical conditions for which a "5-HT₁-like" receptor agonist is indicated, for example, migraine. The carrier must be pharmaceutically acceptable to the recipient and must be compatible with, i.e. not have a deleterious effect upon, the other ingredients in the composition. The carrier may be a solid or liquid and is preferably formulated with at least one compound of formula (I) or (Ia) as a unit dose formulation, for example, a tablet which may contain from 0.05 to 95% by weight of the active ingredient. If desired, other physiologically active ingredients may also be incorporated in the pharmaceutical compositions of the invention.

Possible formulations include those suitable for oral, sublingual, buccal, parenteral (for example, subcutaneous, intramuscular, or intravenous), rectal, topical and intranasal administration. The most suitable means of administration for a particular patient will depend on the nature and severity of the condition being treated and on the nature of the active compound, but, where possible, oral administration is preferred.

Formulations suitable for oral administration may be provided as discrete units, such as tablets, capsules, cachets, or lozenges, each containing a predetermined amount of the

active compound; as powders or granules; as solutions or suspensions in aqueous or non-aqueous liquids; or as oil-in-water or water-in-oil emulsions.

Formulations suitable for sublingual or buccal administration include lozenges comprising the active compound and, typically, a flavoured base, such as sugar and acacia or tragacanth, and pastilles comprising the active compound in an inert base, such as gelatin and glycerin or sucrose and acacia.

Formulations suitable for parenteral administration typically comprise sterile aqueous solutions containing a predetermined concentration of the active compound; the solution is preferably isotonic with the blood of the intended recipient. Although such solutions are preferably administered intravenously, they may also be administered by subcutaneous or intramuscular injection.

Formulations suitable for rectal administration are preferably provided as unit-dose suppositories comprising the active ingredient and one or more solid carriers forming the suppository base, for example, cocoa butter.

Formulations suitable for topical or intranasal application include ointments, creams, lotions, pastes, gels, sprays, aerosols and oils. Suitable carriers for such formulations include petroleum jelly, lanolin, polyethylene glycols, alcohols, and combinations thereof. The active ingredient is typically present in such formulations at a concentration of from 0.1 to 15% w/w.

The formulations of the invention may be prepared by any suitable method, typically by uniformly and intimately admixing the active compound(s) with liquids or finely divided solid carriers, or both, in the required proportions and then, if necessary, shaping the resulting mixture into the desired shape.


For example, a tablet may be prepared by compressing an intimate mixture comprising a powder or granules of the active ingredient and one or more optional ingredients, such as a binder, lubricant, inert diluent, or surface active dispersing agent, or by moulding an intimate mixture of powdered active ingredient and inert liquid diluent.

Aqueous solutions for parenteral administration are typically prepared by dissolving the active compound in sufficient water to give the desired concentration and then rendering the resulting solution sterile and isotonic.


Thus, according to a fifth aspect of the present invention, there is provided the use of a compound of formula (I) or (Ia) in the preparation of a medicament for the prophylaxis or treatment of a clinical condition for which a "5-HT₁-like" receptor agonist is indicated, for example, migraine.

According to a sixth aspect, there is provided a method for the prophylaxis or treatment of a clinical condition in a mammal, for example, a human, for which a "5-HT₁-like" receptor agonist is indicated, for example, migraine, which comprises the administration to said mammal of a therapeutically effective amount of a compound of formula (I) or (Ia) or of a physiologically acceptable salt, solvate, or physiologically functional derivative thereof.

The present invention also provides, in a seventh aspect, a process for the preparation of a compound of formula (I) or (Ia). This process comprises reaction of a compound of formula (II).




wherein W, Z and n are as hereinbefore defined, with a compound of formula (III)



wherein A is as hereinbefore defined.

The reaction is preferably carried out in the presence of a base such as an organic base, for example triethylamine, pyridine or Hunig's base, or an inorganic base, for example sodium hydride or potassium carbonate. Sodium hydride is preferred. The reaction may be carried out in polar aprotic solvents such as dimethylformamide, dimethylsulphoxide or in chlorinated solvents such as dichloromethane. Dimethylformamide is the preferred solvent. The reaction may be carried out in the temperature range 0-100°. 0° to room temperature is preferred.


Where it is desired to prepare a compound of formula (Ia), a compound of formula (IIa)



wherein X, Y, R, n, R^1 and R^2 are as hereinbefore defined may be used in place of the compound of formula (II).

Compounds of formula (II) and (IIa) can be made according to the methods described in EP.A-0486666, incorporated herein by reference. Compounds of formula (III) are commercially available or can be readily prepared by those skilled in the art using known methods.

Alternatively, compounds of formula (I) or (Ia) may be prepared by reaction of a compound of formula (II) or (IIa) with a compound of formula (IV)



wherein A is as hereinbefore defined.

The reaction is preferably carried out in the presence of a base such as an organic base, for example triethylamine, pyridine or dimethylamino pyridine or an inorganic base for example sodium hydride or potassium carbonate. Dimethylaminopyridine is preferred when it is desired to make the compound 4-[1-t-Butyloxycarbonyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl] oxazolidin-2-one acetate. Sodium hydride is the preferred base when it is desired to make the compound 4-[1-Acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate. The reaction may be carried out in polar aprotic solvents such as acetonitrile, dimethylformamide, dimethylsulphoxide or in chlorinated solvents such as dichloromethane. Acetonitrile is the preferred solvent for synthesis of the compound 4-[1-t-Butyloxycarbonyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate. Dimethylformamide is the preferred solvent for synthesis of the compound 4-[1-Acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl] oxazolidin-2-

one acetate. The reaction may be carried out in the temperature range 0-100°. 0° to room temperature is preferred.

Compounds of formula (IV) are commercially available or can be readily prepared by those skilled in the art using known methods.

Salts, solvates and physiologically functional derivatives of Compounds of formula (I) or (Ia) may be prepared from compounds of formula (I) or (Ia) using standard techniques known in the art.

The invention will now be described, by way of illustration only, by the following examples:

Example 1: 4-[1-Benzoyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl] oxazolidin-2-one acetate

4-{1-H-3-[2-(Dimethylamino)ethyl]indol-5-ylmethyl} oxazolidin-2-one (1g, 3.49 mmol) in anhydrous dimethylformamide (DMF) was added by syringe to a suspension of hexane washed sodium hydride (132mg of 60% dispersion, 3.30mmol) in anhydrous DMF (5ml) at room temperature. The gel-like slurry was stirred at room temperature for 1 hour, benzoyl chloride (405µl, 3.49 mmol) was added and the reaction stirred for a further hour. Sodium hydride (132 mg of 60% dispersion) added and the whole stirred overnight at room temperature. The reaction mixture was partitioned between ethyl acetate and saturated aqueous ammonium chloride. The organic phase was washed with brine and dried over magnesium sulphate. The crude product was purified by (a) chromatography on silica, eluting with dichloromethane/methanol/ammonia (90:10:0.5) and then (b) Reverse phase HPLC on a C⁸ column eluting with acetonitrile/water/ammonium acetate to give, after freeze-drying, the title product as a colourless amorphous solid (with 1.0 ACOH).

¹H NMR (200 MHz, d₆ DMSO); 8.26 (1H, d, J = 9Hz); 7.54-7.80 (7H, m); 7.26 (1H, d, J=9Hz); 7.22 (1H, s); 4.28 (1H, t, J = 7.5 Hz); 4.08 (2H, m); 2.92 (2H, m); 2.79 (2H, t); 2.53 (2H, t); 2.18 (6H, s).

The following compounds of examples 2-4 were prepared using a method analogous to that described in example 1:

Example 2: 4-[1-Pivaloyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl] oxazolidin-2-one acetate

-10-

Amorphous colourless solid (with 1.35 mol ACOH)

^1H NMR (200 MHz, d_6 DMSO); 8.28 (1H, d, $J = 9\text{Hz}$); 7.89 (1H, s); 7.78 (1H, s); 7.47 (1H, s); 7.18 (1H, d, $J = 9\text{Hz}$); 4.28 (1H, t, $J = 7.5\text{Hz}$); 4.07 (2H, m); 2.89 (2H, m); 2.86 (2H, t); 2.58 (2H, t); 2.25 (6H, s); 1.46 (9H, s).

Example 3: 4-[1-(2'-toluoyl)-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate

Amorphous colourless solid mp 51.5-54° (with 0.7 ACOH)

^1H NMR (200 MHz, d_6 DMSO); 8.05 (1H, d, $J = 9\text{Hz}$); 7.35-7.59 (5H, m); 7.25 (1H, d, $J = 9\text{Hz}$); 6.94 (1H, s); 4.28 (1H, t, $J = 7.5\text{Hz}$); 4.09 (2H, m); 2.92 (2H, m); 2.77 (2H, t); 2.56 (2H, t); 2.24 (6H, s); 2.18 (3H, s).

Calc. for $\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}_3 \cdot 0.25 \text{H}_2\text{O} \cdot 0.68 \text{ACOH}$; C:65.73, H:6.80, N:8.83.

Found C: 65.68, H:6.89, N:9.08.

Accurate mass; 405.20305 ($\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}_3$)

Example 4: 4-[1-Benzoyloxycarbonyl-3-[2-(dimethylamino)ethyl] indol-5-ylmethyl]oxazolidin-2-one

Amorphous colourless solid

^1H NMR (200 MHz, d_6 DMSO); 7.97 (1H, d, 9Hz); 7.78 (1H, s); 7.38-7.57 (7H, m); 7.21 (1H, d, $J = 9\text{Hz}$); 5.46 (2H, s); 4.27 (1H, t, $J = 7.5\text{Hz}$); 4.07 (2H, m); 2.88 (2H, m); 2.70 (2H, t); 2.56 (2H, t); 2.22 (6H, s).

Example 5: 4-[1-t-Butyloxycarbonyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate

4-{1-H-3-[2-(Dimethylamino)ethyl]indol-5-ylmethyl}oxazolidin-2-one (1.5g, 5.23 mmol) and 4-dimethylaminopyridine (61mg) dissolved in anhydrous acetonitrile (20ml) and dry DMF (5ml). Di-t-butyl dicarbonate (1.25g, 5.75mmol) in dry acetonitrile (10ml) was added, with stirring over 1.5 hours. The reaction mixture was concentrated in vacuo and partitioned between water and ethyl acetate. Organic phase washed with water and dried and residue purified by (a) chromatography on silica, eluting with dichloromethane/methanol/ammonia (90:10:1) and then (b) Reverse phase HPLC on a C^8

-11-

column eluting with acetonitrile/water/ammonium acetate to give after freeze-drying the title compound as colourless amorphous solid (with 0.6 ACOH)

^1H NMR (200 MHz, d_6 , DMSO); 7.96 (1H, d, $J = 9.5$ Hz); 7.77 (1H, s); 7.48 (1H, s); 7.46 (1H, d); 7.20 (1H, d, $J = 9.5$ Hz); 4.27 (1H, t); 4.06 (2H, m); 2.88 (2H, m); 2.78 (2H, t); 2.55 (2H, t); 2.23 (3H, s); 1.53 (9H, s).

Accurate mass; 387.2136 ($\text{C}_{21}\text{H}_{29}\text{N}_3\text{O}_4$)

Example 6: 4-[1-Acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate

4-{1-H-3-[-2-(Dimethylamino)ethyl]indol-5-ylmethyl} oxazolidin-2-one (1g, 3.48 mmol) in anhydrous dimethylformamide (DMF) was added by syringe to a suspension of sodium hydride (139 mg of 60% dispersion, 3.48 mmol) in anhydrous DMF (20ml) at room temperature under nitrogen. The reaction mixture was stirred at room temperature for 2 hours and then cooled to 5°C. Acetic anhydride (0.355g, 3.48mmol) in anhydrous tetrahydrofuran (2ml) was added to the reaction mixture dropwise over 10 minutes keeping the temperature at 5°C stirring continued at this temperature for one hour. The reaction mixture was partitioned between chloroform and saturated aqueous ammonium chloride. The organic phase was dried over magnesium sulphate and the crude product purified by Reverse Phase HPLC on a C8 column eluting with acetonitrile/water/ammonium acetate. Freeze drying afforded the title compound.

^1H NMR (200 MHz, d_6 DMSO); 8.20 (1H, d); 7.77 (1H, s); 7.62 (1H, s); 7.46 (1H, d); 7.18 (1H, d); 4.28 (1H, t); 4.08 (2H, m); 2.90 (2H, m); 2.79 (2H, t); 2.57 (3H, s); 2.53 (2H, t); 2.25 (6H, s); 1.88 (3H, s)

Calc. for $\text{C}_{18}\text{H}_{23}\text{N}_3\text{O}_3 \cdot \text{CH}_3\text{COOH} \cdot 1.09 \text{H}_2\text{O}$; C, 58.73; H, 7.14; N, 10.27.

Found C, 58.62; H, 7.28; N, 10.38.

PHARMACEUTICAL FORMULATION EXAMPLES

In the following Examples, the "active ingredient" may be any compound of formula (I) or (Ia) and/or a physiologically acceptable salt, solvate, or physiologically functional derivative thereof.

(1) Tablet formulations(i) Oral

	<u>Mg/tablet</u>		
	<u>A</u>	<u>B</u>	<u>C</u>
Active ingredient	25	25	25
Avicel	13	-	7
Lactose	78	47	-
Starch (maize)	-	9	-
Starch (pregelatinised, NF15)	-	-	32
Sodium starch glycollate	5	-	-
Povidone	3	3	-
Magnesium stearate	1	1	1
	<hr/>	<hr/>	
	125	85	65

(ii) Sublingual

	<u>Mg/tablet</u>	
	<u>D</u>	<u>E</u>
Active ingredient	25	25
Avicel	10	-
Lactose	-	36
Mannitol	51	57
Sucrose	-	3
Acacia -		3
Povidone	3	-
Magnesium stearate	1	1
	<hr/>	<hr/>
	90	125

-13-

Formulations A to E may be prepared by wet granulation of the first six ingredients with the povidone, followed by addition of the magnesium stearate and compression.

(iii) Buccal

	<u>Mg/tablet</u>
Active ingredient	25
Hydroxypropylmethyl cellulose (HPMC)	25
Polycarbophil	39
Magnesium stearate	1
	<hr/>
	90

The formulation may be prepared by direct compression of the admixed ingredients.

(2) Capsule formulations(i) Powder

	<u>Mg/Capsule</u>	
	<u>F</u>	<u>G</u>
Active ingredient	25	25
Avicel	45	-
Lactose	153	-
Starch (1500 NF)	-	117
Sodium starch glycollate	-	6
Magnesium stearate	2	2
	<hr/>	<hr/>
	225	150

Formulations F and G may be prepared by admixing the ingredients and filling two-part hard gelatin capsules with the resulting mixture.

(ii) Liquid fill

	<u>Mg/Capsule</u>	
	<u>H</u>	<u>I</u>
Active ingredient	25	25
Macrogol 4000 BP	200	-
Lecithin	-	100

-14-

Arachis oil	-	100
	—	—
	225	225

Formulation H may be prepared by melting the Macrogol 4000 BP, dispersing the active ingredient in the melt and filling two-part hard gelatin capsules therewith. Formulation I may be prepared by dispersing the active ingredient in the lecithin and arachis oil and filling soft, elastic gelatin capsules with the dispersion.

(iii) Controlled release

	<u>Mg/tablet</u>
Active ingredient	25
Avicel	123
Lactose	62
Triethylcitrate	3
Ethyl cellulose	12
	—
	225

The formulation may be prepared by mixing and extruding the first four ingredients and spheronising and drying the extrudate. The dried pellets are coated with ethyl cellulose as a release controlling membrane and filled into two-part, hard gelatin capsules.

(3) Intravenous injection formulation

	<u>% by weight</u>
Active ingredient	2%
Hydrochloric acid)	q.s to pH 7
Citrate buffer)	
Water for Injections	to 100%

The active ingredient is taken up in the citrate buffer and sufficient hydrochloric acid added to affect solution and adjust the pH to 7. The resulting solution is made up to volume and filtered through a micropore filter into sterile glass vials which are sealed and oversealed.

BIOLOGICAL EXAMPLES

a) Investigation of Prodrug Activity in Rats of Compound of Example 6

Investigations were carried out in rats to determine whether 4-[1-Acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate (compound of Example 6) acts as a prodrug for the pharmacologically active compound 4-[3-[2-(dimethylamino)ethyl]-1H-indol-5-ylmethyl]-2-oxazolidinone described in EP-A-0486666. The compound of Example 6 was incubated with rat plasma at a concentration of approx. 0.2 µg/ml for up to 4 h. At selected times the incubations were terminated by addition of acetone and the concentrations of both 4-[1-Acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate and its breakdown product 4-[3-[2-(dimethylamino)ethyl]-1H-indol-5-ylmethyl]-2-oxazolidinone were determined by HPLC assay. In addition rats were administered 4-[1-Acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate intravenously at a dose of 4 mg/kg and the plasma and brain concentrations of this compound and the breakdown product, 4-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-ylmethyl]-2-oxazolidinone were determined up to 8 h after administration.

4-[1-Acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate was converted to 4-[3-[2-(Dimethylamino)ethyl]-1H-indol-5-ylmethyl]-2-oxazolidinone in rat plasma, with a half-life of approximately 3 h. After 4 h incubation only 30% of the parent compound of example 6 remained.

After intravenous administration, the peak plasma concentration (C_{max}) of 4-[1-acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate was approx 0.2 µg/ml (5 min) and this parent compound was detectable up to 1 h post dose (limit of detection 0.02 µg/ml); the plasma half-life was approximately 0.2 h. The breakdown product, 4-[3-[2-(dimethylamino)ethyl]-1H-indol-5-ylmethyl]-2-oxazolidinone was also detected in the plasma (C_{max} 0.3 µg/ml, 5 min) up to 4 h after dosing.

These experiments demonstrate that 4-[1-acetyl-3-[2-(dimethylamino)ethyl]indol-5-ylmethyl]oxazolidin-2-one acetate is converted into 4-[3-[2-(dimethylamino)ethyl]-1H-indol-5-ylmethyl]-2-oxazolidinone in rats.

-16-

b) Investigation of Prodrug Activity of Compound of Example 3

The compound of Example 3 was incubated with rat hepaticocytes to determine whether the active compound, 4-[3-[2-(dimethylamino)ethyl]-1H-indol-5-ylmethyl]-2-oxazolidinone, is a main product of its metabolism. After 30 minutes incubation, no compound of Example 3 remained, while approximately 50% had been converted to the active compound.

These results indicate that the compound of Example 3 is extensively and rapidly converted into the active compound.

c) Investigation of Trans-Cellular Absorption of Compound of Example 3

The diffusion rates of the Compound of Example 3 and the active compound, 4-[3-[2-(dimethylamino)ethyl]-1H-indol-5-ylmethyl]-2-oxazolidinone, through a lipoidal membrane barrier (laceryl alcohol/caprylic acid lipid membrane) were determined and compared. The results were :-


Diffusion Rate (cm/min/1000)

Compound of Example 3	3.88
Active compound	0.68

These results indicate that the Compound of Example 3 may be significantly more efficiently absorbed via transcellular mechanisms than the active compound of which it is a pro-drug.

CLAIMS

- 1) A compound of formula (I)




wherein


A is C₁₋₆ alkyl, -O-C₁₋₆ alkyl, -O-phenyl or phenyl, optionally substituted by C₁₋₃ alkyl or halogen;

n is an integer of from 0 to 3;


W is a group of formula (i), (ii) or (iii)



(i)




(ii)



(iii)

wherein R is hydrogen or C₁₋₄ alkyl, x is -O-, -S-, -NH-, or -CH₂-, Y is oxygen or sulphur and the chiral centre * in formula (i) or (ii) is in its (S) or (R) form or is a mixture thereof in any proportions; and


Z is a group of formula (iv), (v) or (vi)



(iv)



(v)




(vi)

wherein R¹ and R² are independently selected from hydrogen and C₁₋₄ alkyl and R³ is hydrogen or C₁₋₄ alkyl;

and salts, solvates and physiologically functional derivatives thereof.

-18-

2. A compound of formula (Ia)



wherein

A is C₁₋₆ alkyl, -O-C₁₋₆ alkyl, -O-phenyl or phenyl, optionally substituted by C₁₋₃ alkyl or halogen;

n is an integer of from 0 to 3;

R is hydrogen or C₁₋₄ alkyl;

X is -O-, -S-, -NH- or -CH₂-;


Y is oxygen or sulphur; and

R¹ and R² are independently selected from hydrogen and C₁₋₄ alkyl, and salts, solvates, and physiologically functional derivatives thereof.


3. A compound according to claim 1 or claim 2 wherein R¹ and R² are both methyl, and salts, solvates and physiologically functional derivatives thereof.
4. A compound according to claim 2 or claim 3 wherein X is -O- and Y is oxygen, and salts, solvates and physiologically functional derivatives thereof.
5. A pharmaceutical formulation comprising at least one compound of formula (I) or formula (Ia) as defined herein together with one or more pharmaceutically acceptable carriers and/or excipients.
6. A compound of formula (I) or formula (Ia) as defined herein for use in therapy.
7. The use of a compound of formula (I) or (Ia) as defined herein in the manufacture of a medicament for use in the prophylaxis or treatment of migraine.
8. A process for the preparation of a compound according to any of claims 1 to 6 which process comprises

-19-

a) reaction of a compound of formula (II)




wherein W, n and Z are as defined in claim 1 with a compound of formula (III)



wherein A is as defined in claim 1


or

b) reaction of a compound of formula (II) with a compound of formula (IV)



optionally followed by salt, solvate or derivative formulaion.

9. A process according to claim 8 for the preparation of a compound of formula (Ia) which process comprises reaction of a compound of formula (IIa)



with a compound of formula (III) or formula (IV).

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 95/01249

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D413/06 A61K31/405

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,91 18897 (THE WELLCOME FOUNDATION LTD.) 12 December 1991 cited in the application see claims ---	1,7
A	EP,A,0 313 397 (THE WELLCOME FOUNDATION LTD.) 26 April 1989 cited in the application see claims ---	1,7
A	WO,A,93 21182 (MERCK SHARP & DOHME LTD.) 28 October 1993 * page 6, line 1-9; claims * ---	1,7
A	WO,A,92 13856 (PFIZER INC.) 20 August 1992 * example 8A(b), 8B(b), 19, 20; page 13, line 1-5; claims * ---	1,7
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 July 1995

Date of mailing of the international search report

27. 07. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Int ional Application No
PCT/GB 95/01249

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	EP,A,0 610 134 (ADIR ET COMPAGNIE) 10 August 1994 * claims, R2= ...acyle, * -----	1,7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/01249

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9118897	12-12-91	AU-B- 646871	10-03-94
		AU-A- 7957091	31-12-91
		EP-A- 0486666	27-05-92
		EP-A- 0636623	01-02-95
		JP-T- 5502679	13-05-93
		NZ-A- 238424	23-12-93
		PL-B- 166214	28-04-95
		US-A- 5399574	21-03-95
EP-A-313397	26-04-89	AU-A- 2418188	27-04-89
		DE-A- 3881472	08-07-93
		DE-T- 3881472	16-12-93
		ES-T- 2054825	16-08-94
		JP-A- 1146882	08-06-89
		US-A- 5225431	06-07-93
WO-A-9321182	28-10-93	AU-B- 4076693	18-11-93
		EP-A- 0636131	01-02-95
		JP-T- 7505649	22-06-95
WO-A-9213856	20-08-92	AU-B- 655456	22-12-94
		AU-A- 1263792	07-09-92
		CZ-A- 9301656	18-05-94
		EP-A- 0571471	01-12-93
		HU-A- 65766	28-07-94
		JP-T- 6500122	06-01-94
		NZ-A- 241584	22-12-94
		ZA-A- 9200969	11-08-93
EP-A-610134	10-08-94	FR-A- 2701026	05-08-94
		AU-B- 5481794	04-08-94
		CA-A- 2114593	03-08-94
		JP-A- 6271571	27-09-94
		NZ-A- 250800	28-03-95
		US-A- 5348968	20-09-94